

# On Virtually Projective Groups

P. A. Zalesskii\*

To A.E. Zalesskii on the occasion of his 65-th birthday

## Abstract

It is proved that the quotient  $G/\langle \text{tor}(G) \rangle$  of a virtually projective profinite group  $G$  modulo its normal subgroup generated by all torsion of  $G$  is projective.

2000 Mathematics Subject Classification: 20E18

Key words: profinite groups.

## 0. Introduction

Let  $G$  be a virtually free group. Then by results of Karrass, Magnus, Solitar, Cohen and Scott  $G = \pi_1(\mathcal{G}, \Gamma)$  is the fundamental group of a graph of finite groups  $(\mathcal{G}, \Gamma)$ . By the central result of Bass-Serre's theory of groups acting on trees this is equivalent to the fact that  $G$  acts on a tree  $S$  with finite vertex stabilizers such that  $S/G = \Gamma$ . Let  $\text{tor}(G)$  be the set of all nontrivial torsion elements of  $G$ . Since every torsion element must fix a vertex of  $S$ , it follows that the group  $\langle \text{tor}(G) \rangle$  is generated by the stabilizers of vertices of  $S$  and therefore  $S/\langle \text{tor}(G) \rangle$  is a tree on which  $G/\langle \text{tor}(G) \rangle$  acts freely. Thus  $G/\langle \text{tor}(G) \rangle = \pi_1(\Gamma)$  is the fundamental group of the graph  $\Gamma$  and hence is free.

It was proved recently in [HZ-2002] that a finitely generated virtually free pro- $p$  group is the fundamental pro- $p$  group of a finite graph of finite  $p$ -groups. Unfortunately, this result does not hold in the infinitely generated case. However, there is still hope that a virtually free pro- $p$  group acts on a pro- $p$  tree with finite vertex stabilizers, because in the pro- $p$  case this is a weaker property than to be the fundamental group of a graph of finite  $p$ -groups. Moreover, it is shown in [HZ-2002] that  $G/\langle \text{tor}(G) \rangle$  is free pro- $p$ , when  $G$  is second countable that would be the consequence of this conjecture if proved in this case.

The situation in the profinite case is more complicated. A virtually free profinite group does not act in general on a profinite tree and so does not have a structure similar to a discrete virtually free group. An example is the semidirect product  $\hat{\mathbf{Z}} \rtimes C_2$ , where  $C_2$  inverts elements of the 2 component  $\mathbf{Z}_2$  of  $\hat{\mathbf{Z}}$  and fixes the elements of  $p$  components  $\mathbf{Z}_p$  for all other primes  $p$ .

The objective of the present paper is to show that nevertheless one can obtain quite reasonable information on  $G/\langle \text{tor}(G) \rangle$  of virtually free profinite group  $G$ . In fact, our result is even more general.

**Theorem** Let  $G$  be a virtually projective profinite group. Then  $G/\langle \text{tor}(G) \rangle$  is projective.

---

\* Supported by CNPq.

In the case  $\langle \text{tor}(G) \rangle = 1$  ( i.e. when  $G$  is torsion free) the result is due to Serre [S-1965]. Note that free groups, free pro- $p$  groups and projective groups are exactly groups of cohomological dimension 1 in the categories of groups, pro- $p$  groups and profinite groups, respectively. Therefore one could ask as a possible generalization of the theorem above whether for a group  $G$  of finite virtual cohomological dimension  $n$  one has  $cd(G/\langle \text{tor}(G) \rangle) \leq n$ . In Section 3 we give an example of a group of virtual cohomological dimension 2 whose quotient  $G/\langle \text{tor}(G) \rangle$  is even not torsion free, and so has infinite cohomological dimension. This shows that the situation with groups of (virtual) cohomological dimension 1 is rather special.

The structure of the paper as follows. In Section 1 the ideas of [RHZ-99] and [HZ-2000] are used to complete the result in the pro- $p$  case. The main result is proved in Section 2.

In Section 3 besides the example mentioned above we also give an example of a semidirect product  $F \rtimes C_2$  of a free pro-2 group  $F$  of uncountable rank and a group of order 2 that does not satisfy the Dyer-Scott type decomposition

$$G = \prod_{x \in X} (H_x \times C_2) \amalg H,$$

where  $H_x$  and  $H$  are free pro-2 groups. When  $F$  is of countable rank the Dyer-Scott decomposition holds (see Theorem 1.2 below).

The necessary material on profinite groups (like a notion of a free profinite group on a topological space) can be found in [RZ-2000] and [W-98]. The definition of a free pro- $p$  product which is used in the paper can be found in [NSW-2000], Chapter IV, S3, or in [M-1989]. We shall use frequently Serre's result from [S-1965] that states that a virtually projective torsion free profinite group is projective.

Notation. All groups in the paper are profinite, homomorphisms are continuous and subgroups are closed. By  $p$  will be denoted usually a prime number. For a pro- $p$  group  $G$  we denote the Frattini subgroup of  $G$  by  $\Phi(G)$ .  $\text{tor}(G)$  means the subset of torsion elements of  $G$  and  $x^g$  stands for  $x^{-1}gx$ . For a profinite space  $X = \varprojlim X_i$ ,  $|X_i| < \infty$  and a profinite ring  $R$  we denote by  $[[RX]] = \varprojlim [RX_i]$  a free profinite module over the space  $X$ .

## 1. The pro- $p$ case

Denote by  $\mathbf{n}(G)$  the index of maximal free pro- $p$  normal subgroup of  $G$ . The proof of the result in this case uses induction on  $\mathbf{n}(G)$ . We first formulate a theorem that gives the base of induction.

Let  $G$  be a pro- $p$  group having an open free pro- $p$  subgroup  $F$ . Then the set  $\mathcal{T}$  of all subgroups of order  $p$  is a profinite space, since it is a projective limit of corresponding finite discrete spaces of quotients  $G/U$ , where  $U$  runs through the all open normal subgroups of  $G$  which are contained in  $F$ . Moreover,  $G$  acts continuously on  $\mathcal{T}$  by conjugation.

The stabilizer  $G_T$  of  $T \in \mathcal{T}$  with respect to this action is just the centralizer  $G_T = C_G(T)$ . We denote by  $\varphi_G : \mathcal{T} \rightarrow \mathcal{T}/G$  the natural map. Then for  $t \in \mathcal{T}/G$  the preimage

$\varphi_G^{-1}(t)$  is the  $G$ -orbit. The next lemma is just a homological version of the result of Scheiderer from [Sch-1994, Theorem 12.13].

**Lemma 1.1.** (Lemma 5 in [HRZ-1999]) *For any  $n \geq 2$  the canonical homomorphism*

$$\varphi_n : \bigoplus_{t \in \mathcal{T}/G} H_n(G, [\mathbf{F}_p \varphi_G^{-1}(t)]) \rightarrow H_n(G, \mathbf{F}_p)$$

*is a topological isomorphism.*

Now we state a pro- $p$  version of the Dyer-Scott theorem [DS-1975] that was proved in [Sch-1999] for finitely generated case and in [HRZ-1999] in the form below. Note that the result holds upon the condition of the existence of a continuous section  $\mathcal{T}/G \rightarrow \mathcal{T}$ . Proposition 3.3 shows that this condition is essential.

**Theorem 1.2.** (Proposition 9 in [HRZ-1999]) *Let  $G$  be a pro- $p$  group having free subgroup  $F$  of index  $p$ . Suppose there exists a continuous section  $\sigma : \mathcal{T}/G \rightarrow \mathcal{T}$ . Put  $T = \sigma(t)$  regarding as a subgroup of  $G$ . Then*

$$G = \left( \prod_{T \in \text{im}(\sigma)} (T \times C_F(T)) \right) \amalg H,$$

*is a free pro- $p$  product over the profinite space  $\mathcal{T}/G$ , where  $H$  is a free pro- $p$  subgroup of  $F$ .*

We note that a section  $\sigma$  always exists if the action is free or if  $\mathcal{T}$  is second countable (see Lemmas 5.6.5 and 5.6.7 in [RZ-2000]).

The next proposition is extracted from the proof of Proposition 13 in [HRZ-1999]

**Proposition 1.3.** *Let  $G$  be a pro- $p$  group having a free pro- $p$  subgroup  $F$  of index  $p$ . Then  $G$  embeds into a free pro- $p$  product*

$$G_0 = (C_p \times H) \amalg H_0 \tag{6}$$

*where  $H, H_0$  are free pro- $p$  groups and  $C_p$  is a group of order  $p$ .*

*Proof:* If  $G$  is free pro- $p$ , there is nothing to prove. So assume that  $G$  is not free pro- $p$ ; then by Serre's result the torsion  $\text{tor}(G) \neq \emptyset$ . Let  $\varphi : G \rightarrow G/F$  be the natural epimorphism. Choose a generator  $c$  of  $G/F \cong C_p$  and put  $C = \text{tor}(G) \cap \varphi^{-1}(c)$ . For  $T \in \mathcal{T}$  denote by  $c_T$  the unique element of  $C \cap T$ .

For the rest of the proof fix an arbitrary  $T_0 \in \mathcal{T}$  and write  $c_0 = c_{T_0}$ . The set

$$c_0^{-1}C \subset F \tag{2}$$

is naturally homeomorphic to the pointed boolean space  $(\mathcal{T}, T_0)$ . This way, in the sequel,  $(\mathcal{T}, T_0)$  will appear as an indexing profinite space. Let  $F(\mathcal{T}, T_0)$  be a free pro- $p$  group over

the pointed space  $(\mathcal{T}, T_0)$ . We shall denote by  $z_T$  the image of a point  $T \in \mathcal{T}$  under the natural injection  $\mathcal{T} \longrightarrow F(\mathcal{T}, T_0)$ . Form the free pro- $p$  product

$$F_0 = F(\mathcal{T}, T_0) \amalg F. \quad (3)$$

Observe that  $c_0^{-1}c_T \in F$  for every  $T \in \mathcal{T}$ . Define an automorphism  $\alpha_0 \in \text{Aut}(F_0)$  by putting

$$\begin{aligned} \alpha_0(z_T) &= c_0^{-1}c_T z_T & T \in \mathcal{T}, \\ \alpha_0(f) &= c_0^{-1}f c_0 & f \in F. \end{aligned} \quad (4)$$

We check that  $\alpha_0$  has order  $p$  by looking at generators of  $F_0$ . We show first by induction on  $k$  that

$$\alpha_0^k(z_T) = c_0^{-k}c_T^k z_T$$

for all  $T \in (\mathcal{T}, T_0)$  and  $1 \leq k \leq p$ . The formula follows from the definition of  $\alpha_0$  for  $k = 1$ . Assuming that the formula holds for  $k - 1$ , one has

$$\begin{aligned} \alpha_0^k(z_T) &= \alpha_0^{k-1}\alpha_0(z_T) = \alpha_0^{k-1}(c_0^{-1}c_T z_T) = c_0^{-k+1}c_0^{-1}c_T c_0^{k-1}\alpha_0^{k-1}(z_T) \\ &= c_0^{-k}c_T c_0^{k-1}c_0^{-k+1}c_T^{k-1}z_T = c_0^{-k}c_T^k z_T \end{aligned}$$

as required.

Hence  $\alpha_0^p(z_T) = z_T$  and certainly,  $\alpha_0^p(f) = f c_0^p = f$  for any  $f \in F$ .

There is a natural embedding of  $G$  into  $G_0 := F_0 \rtimes \langle \alpha_0 \rangle$  where  $F$  is sent to a copy of  $F$  in  $G_0$  and  $c_0$  is sent to  $\alpha_0$ . We shall identify  $\alpha_0$  and  $c_0$  henceforth.

By construction, the torsion of  $G_0$  coincides with the torsion of  $G$ , and since, as a simple consequence of Eq.(4) and the identification  $\alpha_0 = c_0$

$$z_T c_0 z_T^{-1} = c_T \quad (5)$$

holds for  $T \in \mathcal{T}$ ,  $G_0$  has only one conjugacy class of subgroups of order  $p$ . An application of Theorem 1.2 then yields a decomposition

$$G_0 = (C_p \times H) \amalg H_0 \quad (6)$$

with  $H, H_0$  suitable free pro- $p$  groups of  $F_0$ . □

**Corollary 1.4.** *Let  $G$  be a pro- $p$  group having a free pro- $p$  subgroup  $F$  of index  $p$ . Then*

- (i)  $G/\langle \text{tor}(G) \rangle$  is free pro- $p$ ;
- (ii)  $C_F(c)$  is a free factor of  $F$  for any torsion element  $c$  of  $G$ .
- (iii) If  $G$  is generated by torsion, then there exists a continuous section  $\sigma : T/G \longrightarrow T$  and one has

$$G = \left( \prod_{T \in \text{im}(\sigma)} T \right).$$

*Proof:* (i) By the preceding theorem  $G$  embeds into a free pro- $p$  product

$$G_0 = (C_p \times H) \amalg H_0 \quad (6)$$

where  $H, H_0$  are free pro- $p$  groups. Let  $X$  and  $X_0$  be closed bases of  $H$  and  $H_0$  respectively. Hence  $G_0$  can be viewed as an HNN-group  $\langle C_p, X, X_0 \mid xcx^{-1} = c \text{ for } c \in C_p, x \in X \rangle$ . It follows that  $G_0$  acts on a pro- $p$  tree  $S$  whose vertex stabilizers are conjugates of  $C_p$  (see Proposition 3.8 in [ZM-1990]). Then  $G$  acts on  $S$  as well and  $\langle \text{tor}(G) \rangle$  is exactly the subgroup of  $G$  generated by the vertex stabilizers. So by Corollary 3.6 in [RZ1-2000]  $G/\text{tor}(G)$  is free pro- $p$  as required.

(ii) Since every torsion element is conjugate in  $G_0$  to some element of  $C_p$ , using conjugation if necessary, we may assume that  $C_p = \langle c \rangle$ . Let  $f : G_0 \rightarrow H$  be the epimorphism that sends  $C_p$  and  $H_0$  to 1 and  $H$  identically onto  $H$ . The restriction of  $f$  to  $C_F(c)$  is injective, because  $C_{G_0}(C_p) = C_p \times H$  (see Corollary 4 in [RZ1-2000]). Hence  $F$  splits as a semidirect product  $F = M \rtimes C_F(c)$ . It follows that  $\Phi(F) \cap C_F(c) = \Phi(C_F(c))$ . Then by Lemma 9.1.18 in [RZ-2000]  $C_F(c)$  is a free factor of  $F$ .

(iii) Let  $T$  be a subgroup of  $G$  of order  $p$ . Conjugating it if necessary we may assume that  $T = C_p$ . Let  $\varphi : G_0 \rightarrow H \amalg H_0$  be the epimorphism that sends  $C_p$  to 1 and  $H, H_0$  identically to their copies in  $H \amalg H_0$ . As  $C_{G_0}(C_p) = C_p \times H$  (see Corollary 4 in [RZ1-2000]), the restriction of  $\varphi$  to  $C_F(T)$  is injective. Since  $G$  is generated by torsion and every torsion element is conjugate in  $G_0$  to some element of  $C_p$  (cf. Theorem 4.2 (a) in [RZ1-2000]), one has  $\varphi(G) = 1$ . Hence  $C_F(T) = 1$  for any subgroup  $T$  of  $G$  of order  $p$ . It follows that  $F$  acts freely on  $\mathcal{T}$  and so there exists a section  $\sigma : \mathcal{T}/F \rightarrow \mathcal{T}$  (Lemma 5.6.5 in [RZ-2000]). But  $\mathcal{T}/G = \mathcal{T}/F$ , so the result follows from Theorem 1.2.  $\square$

A finitely generated version of the next theorem is due to Scheiderer [Sch-1999].

**Theorem 1.5.** *Suppose  $F$  is a free pro- $p$  group and  $P$  is a finite  $p$ -group of automorphisms of  $F$ . Then the set of fixed points  $C_F(P)$  is a free factor of  $F$ . In particular, if the rank of  $F$  is finite, so is rank of  $C_F(P)$ .*

*Proof.* Let  $P$  be a nontrivial finite  $p$ -group of automorphisms of  $F$  of minimal order such that the theorem fails. Consider the holomorph  $G = F \rtimes P$ . By Corollary 1.4 (ii),  $|P| > p$ . Pick an element  $c$  in the center of  $P$  with  $c^p = 1$ . By the above case  $C_F(c)$  is a free factor of  $F$ . Therefore  $P/\langle c \rangle$  acts on  $C_F(c)$ , and from the minimality assumption we conclude the result.  $\square$

**Remark 1.6** If  $\alpha$  is an automorphism of order  $p^\infty$  of a finitely generated free pro- $p$  group  $F$ , then it is not known whether the subgroup of fixed point  $C_F(\alpha)$  is finitely generated.

**Proposition 1.7.** *Let  $G$  be any virtually free pro- $p$  group and  $N \triangleleft G$  a normal subgroup of  $G$  generated by torsion elements. Then the following statements hold:*

- (i)  $\text{tor}(G/N) = \text{tor}(G)N/N$  (torsion from  $G/N$  can be lifted).
- (ii)  $G/\langle \text{tor}(G) \rangle$  is free pro- $p$ .

*Proof:*

**Claim 1** (i) and (ii) are equivalent.

For showing (i)  $\Rightarrow$  (ii) pick  $\bar{g} \in G/\langle \text{tor}(G) \rangle$  with  $\bar{g}^p = 1$ . Apply (i) with  $N := \langle \text{tor}(G) \rangle$ , in order to find  $x \in \text{tor}(G)$  with  $x\langle \text{tor}(G) \rangle/\langle \text{tor}(G) \rangle = \bar{g}$ . Since  $x \in \langle \text{tor}(G) \rangle$  conclude  $\bar{g} = 1$ . So  $G/\langle \text{tor}(G) \rangle$  is torsion free. To show that it is free pro- $p$  we use induction on  $\mathbf{n}(G)$ . Let  $c$  be a central element of  $G/F$  of order  $p$ . Then the preimage  $G_1$  of  $\langle c \rangle$  in  $G$  satisfies the assumption of Corollary 1.4 and so  $G_1/\langle \text{tor}(G_1) \rangle$  is free pro- $p$ . Now from (i)  $\text{tor}(G)\langle \text{tor}(G_1) \rangle/\langle \text{tor}(G_1) \rangle = \text{tor}(G/G_1)$  and  $\mathbf{n}(G/G_1) < \mathbf{n}(G)$ . So from the induction hypothesis we deduce that  $G/\langle \text{tor}(G) \rangle = (G/G_1)/\langle \text{tor}(G/G_1) \rangle$  is free pro- $p$  as needed.

Suppose “(ii)  $\Rightarrow$  (i)” is false. Then there exists a virtually free pro- $p$  group  $G$  having a normal subgroup  $N$  generated by torsion and an element  $g \in G$  such that  $gN/N \in \text{tor}(G/N)$  and  $gN \cap \text{tor}(G) = \emptyset$ . Then  $G$  replaced by  $\langle g, N \rangle$  is still a counter example, so we may assume that  $G = \langle g, N \rangle$  and denote such a counter example by  $(g, N)$ . Among the all such counter examples choose one with  $[G : N]$  minimal. We prove first that  $[G : N] = p$ .

Suppose not. Put  $M := \langle g^p, N \rangle$  then  $g^p \notin N$  and  $[M : N] < [G : N]$  so that  $(g^p, N)$  cannot be a counter example. Hence  $M = \langle \text{tor}(M) \rangle$ . On the other hand,  $[G : M] < [G : N]$ , so that  $(g, M)$  is not a counter example either, hence exists  $g_0 \in \text{tor}(G)$  with  $g_0M/M = gM/M$ . Then  $\langle g_0, N \rangle = \langle g, N \rangle$ , i.e.,  $g_0 \in gN \cap \text{tor}(G) = \emptyset$ , a contradiction. Thus  $[G : N] = p$ .

For finishing the proof of Claim 1, note that (ii) implies  $G = \langle \text{tor}(G) \rangle$ . Therefore, there exists a torsion element  $g_0 \in G \setminus N$  and, for suitable  $1 \leq k \leq p - 1$ , one must have  $g_0^k \in gN \cap \text{tor}(G) = \emptyset$ , a contradiction. Therefore Claim 1 is established.

We continue the proof of the proposition. Suppose it is false. Then there exists  $G$  with  $G/\langle \text{tor}(G) \rangle$  not free pro- $p$ . Choose one with  $\mathbf{n}(G)$  minimal and let  $F \triangleleft G$  be a free pro- $p$  group with  $[G : F] = \mathbf{n}(G)$ . Then, in light of Claim 1, there exists  $g \in G$  such that  $g\langle \text{tor}(G) \rangle/\langle \text{tor}(G) \rangle$  of order  $p$  and  $g\langle \text{tor}(G) \rangle \cap \text{tor}(G) = \emptyset$ . It follows that  $\langle g, \text{tor}(G) \rangle$  is still a counter example and since  $[\langle g, \text{tor}(G) \rangle : (\langle g, \text{tor}(G) \rangle \cap F)] \leq \mathbf{n}(G)$  it follows that  $\mathbf{n}(\langle g, \text{tor}(G) \rangle) = \mathbf{n}(G)$ . So from now on we may assume that  $G = \langle g, \text{tor}(G) \rangle$ .

**Claim 2**  $G/F$  is not cyclic.

Suppose it is. Let  $G_i$  be the preimage in  $G$  of the cyclic subgroup of order  $p^i$  in  $G/F$ . Choose  $i$  maximal such that  $G_i = F \rtimes C_{p^i}$ . Then  $\text{tor}(G) \subseteq G_i$ ; indeed, if  $g \in \text{tor}(G) \setminus G_i$  then since  $G/F$  is cyclic,  $g$  has order at least  $p^{i+1}$  and  $G_{i+1} = F \rtimes \langle g \rangle$  contradicting the choice of  $i$ . Put  $n = \mathbf{n}(G)$ . Then it follows from the minimality assumption on  $n$  that  $i = n - 1$  and  $G_{n-1} = \langle \text{tor}(G) \rangle$ . Let  $U$  be a normal subgroup generated by all  $p^{n-2}$  powers of elements of order  $p^{n-1}$ . Then by the minimality assumption on  $n$  and Claim 1 one has  $\text{tor}(G_{n-1}/U) = \text{tor}(G_{n-1})U/U = \text{tor}(G)U/U$ . Consequently as it was shown above  $G_{n-1}/U = F_0 \rtimes C_{p^{n-2}}$  for some free pro- $p$  group  $F_0$  and so  $\mathbf{n}(G_{n-1}/U) = n - 2$ . Therefore it follows from Claim 1 that to use the minimality assumption on  $n$  we have to prove the equality  $G/\langle \text{tor}(G) \rangle = (G/U)/\langle \text{tor}(G/U) \rangle$ .

Suppose not and  $k \in G \setminus \langle \text{tor}(G) \rangle$  such that  $kU/U$  is of finite order. Put  $K = \langle k, U \rangle$  and let  $M$  be a subgroup of  $G$  generated by all elements of order  $p^{n-1}$ . Since any element

of order  $p^{n-1}$  centralizes its  $p^{n-2}$ -power,  $\mathcal{T}_U/M = \mathcal{T}_U/U = \mathcal{T}_U/(F \cap U)$ . Since  $U$  is generated by its torsion Corollary 1.4 (iii) implies that there exists a continuous section  $\sigma_1 : \mathcal{T}_U/U \longrightarrow \mathcal{T}_U$  with

$$U = \coprod_{T \in \text{im}(\sigma_1)} T.$$

Note that the action of  $K/M$  on  $\mathcal{T}_U/M = \mathcal{T}_U/U = \mathcal{T}_U/(F \cap U)$  is free. Indeed, if not then there exists  $A \in \mathcal{T}_U$  and  $f \in F \cap U$  with  $A^k = A^f$ , because  $\mathcal{T}_U/M = \mathcal{T}_U/F \cap K$ ; but then  $kf$  centralizes  $A$  and, since  $C_U(A) = A$  (a free factor is self-centralized, cf. Corollary 4.4 in [RZ1-2000]), the element  $kf$  has to be of finite order; but  $F \cap U \leq \langle \text{tor}(G) \rangle$ , so  $k \in \langle \text{tor}(G) \rangle$ , a contradiction with the choice of  $k$ . Therefore there exists a continuous section  $\sigma : \mathcal{T}/K \longrightarrow \mathcal{T}/U = \mathcal{T}/M$  (cf. Lemma 5.6.5 in [RZ-2000]).

Let  $c$  be a generator of  $K/M$ . Then the  $\mathcal{T}_i := \text{Im}(\sigma)c^i$  ( $i = 0, \dots, p-1$ ) form a partition of  $\mathcal{T}/M = \mathcal{T}/U$  into  $p$  clopen subsets. Define  $K_i := \coprod_{t \in \sigma_1(\mathcal{T}_i)} T$  and write  $\tilde{F} = (F \cap K_i \mid i = 1, \dots, p-1)_K$  to be the normal closure of  $F \cap K_i$ 's. Then  $U = \coprod_{i=0}^{p-1} K_i$ ,

$$U/\tilde{F} = \coprod_{i=0}^{p-1} C_p \tag{*}$$

is a free pro- $p$  product of groups of order  $p$  and the conjugacy classes of the free factors are permuted by the action of  $K/M$ . It follows that the abelianization  $(K/\tilde{F})/(K/\tilde{F})'$  is of order  $p^n$  and since  $F \cap K$  contains the commutator subgroup  $K'$ , the commutator subgroup  $(K/\tilde{F})'$  coincides with  $F \cap K/\tilde{F}$  showing that  $(K/\tilde{F})/(K/\tilde{F})'$  is cyclic of order  $p^n$ . But then  $K/\tilde{F}$  has to be procyclic, a contradiction to (\*).

Thus  $G/\langle \text{tor}(G) \rangle = (G/U)/\langle \text{tor}(G/U) \rangle$  and by the minimality assumption on  $n$  we deduce that Claim 2 holds.

**Claim 3**  $G/F \cong C_p \times C_p$  is the direct product of groups of order  $p$ .

Since  $G/F$  is not cyclic, there exists a normal subgroup  $M$  of  $G$  containing  $F$  such that  $G/M \cong C_p \times C_p$ . For any  $A \subseteq G$  write  $\bar{A} := A\langle \text{tor}(M) \rangle / \langle \text{tor}(M) \rangle$ . Let  $K$  be an arbitrary normal subgroup of index  $p$  in  $G$  containing  $M$ . Then  $\mathbf{n}(K) < \mathbf{n}(G)$  and therefore, by the minimality of  $\mathbf{n}(G)$  and Claim 1,  $\text{tor}(\bar{K}) = \overline{\text{tor}(K)}$ . But every torsion element of  $G$  belongs to some  $K$  of this sort, showing  $\text{tor}(\bar{G}) = \overline{\text{tor}(G)}$  and hence  $\bar{G}/\langle \text{tor}(\bar{G}) \rangle = \bar{G}/\langle \text{tor}(G) \rangle$ . If  $M$  is non-trivial, then  $\mathbf{n}(\bar{G}) < \mathbf{n}(G)$  implying that  $G/\langle \text{tor}(G) \rangle \cong \langle \bar{G}/\text{tor}(\bar{G}) \rangle$  is free pro- $p$ , a contradiction. Hence  $M = 1$  and Claim 3 holds.

Returning to proving the proposition, put  $L := \langle g, F \rangle$ . Then, as  $G/F \cong C_p \times C_p$  deduce  $[G : L] = [G : \langle \text{tor}(G) \rangle] = p$  and  $L \cap \langle \text{tor}(G) \rangle = F$ . On the other hand,  $L$  cannot be torsion free, else, by Serre's result ([S-1965], it is free pro- $p$ , and, being of index  $p$  in  $G$ , the group  $G$  would be free pro- $p$  by cyclic, contradicting Claim 2. Then  $1 \neq \langle \text{tor}(L) \rangle \leq L \cap \langle \text{tor}(G) \rangle = F$  follows, a clear contradiction.  $\square$

## 2. The profinite case

**Lemma 2.1.** *Let  $G$  be a virtually free pro- $p$  group and  $f : G \rightarrow H$  a homomorphism to a profinite group  $H$  that sends every torsion element to 1. Let  $A$  be a discrete  $p$ -primary  $\widehat{\mathbf{Z}}[[H]]$ -module viewed as a  $\widehat{\mathbf{Z}}[[G]]$ -module via  $f$ . Then the induced map  $H^2(H, A) \rightarrow H^2(G, A)$  is the 0-map.*

*Proof:* Clearly  $f$  factors through  $G/\langle \text{tor}(G) \rangle$  and so one has the commutative diagram

$$G[r][rd]G/\langle \text{tor}(G) \rangle[d]H$$

inducing the commutative diagram

$$H^2(G, A)H^2(G/\langle \text{tor}(G) \rangle, A)[l]H^2(H, A)[u][lu]^\varphi$$

By Proposition 1.7  $G/\langle \text{tor}(G) \rangle$  is free pro- $p$  and so  $H^2(G/\langle \text{tor}(G) \rangle, A) = 0$ . It follows that  $\varphi$  is the 0-map.  $\square$

**Lemma 2.2.** *Let  $G$  be a virtually projective group and  $M$  a normal subgroup of  $G$  generated by elements of order coprime to  $p$  and containing all elements of  $G$  of  $p$ -power order. Then the quotient group  $S_p M/M$  is free pro- $p$  for any Sylow  $p$ -subgroup  $S_p$  of  $G$ .*

*Proof:* It suffices to show that  $H^2(G/M, A) = 0$  for any simple  $p$ -primary module  $\widehat{\mathbf{Z}}[[G/M]]$ -module  $A$  (cf. Proposition 7.1.4 and Theorem 7.3.1 in [RZ-2000]). Define the action of  $G$  on  $A$  via  $G/M$ . Consider the 5-term Hochschild-Serre sequence

$$\begin{aligned} 0 \longrightarrow H^1(G/M, A) &\xrightarrow{\text{Inf}} H^1(G, A) \xrightarrow{\text{Res}} \\ H^1(M, A)^{G/M} &\xrightarrow{\text{tr}} H^2(G/M, A) \xrightarrow{\text{Inf}} H^2(G, A) \end{aligned}$$

Since  $M$  is generated by elements of order coprime to  $p$  one has

$$H^1(M, A)^{G/M} = \text{Hom}(M, A)^{(G/M)} = 0.$$

So it remains to prove that  $H^2(G/M, A) \xrightarrow{\text{Inf}} H^2(G, A)$  is the 0-map.

First note that  $H^2(G, A) \rightarrow H^2(S_p, A)$  is an injection (cf. Lemma 10.2.1 in [W-1998]). Moreover, the commutative diagram

$$S_p[r][rd]G[d]G/M$$

induces the commutative diagram

$$H^2(S_p, A)H^2(G, A)[l]H^2(G/M, A)[u][lu]^\varphi$$

and it suffices to show that  $\varphi$  is the 0-map. However, this is the subject of Lemma 2.1.  $\square$

**Corollary 2.3.** *Let  $G$  be a virtually projective group and  $M$  be the normal subgroup generated by all elements of order coprime to  $p$ . Then  $\text{tor}(G)M/M = \text{tor}(G/M)$ .*



*Proof:* For  $g \in G$  with  $gM/M$  of order  $p$  Lemma 2.2 implies that  $\langle g, M \rangle$  must have torsion outside of  $M$  as needed.  $\square$

**Lemma 2.4.** *Let  $G$  be a virtually projective group and  $G/\langle \text{tor}(G) \rangle$  is a pro- $p$  group. Then  $G/\langle \text{tor}(G) \rangle$  is free pro- $p$ .*

*Proof:* Let  $F$  be a maximal open normal projective subgroup of  $G$ . Denote by  $O_p(F)$  the kernel of  $F$  on its maximal pro- $p$  quotient. Note that  $O_p(G) \leq \langle \text{tor}(G) \rangle$ . We show that  $\langle \text{tor}(G) \rangle/O_p(F) = \langle \text{tor}(G/O_p(F)) \rangle$ .

Indeed, let  $s$  be an element of  $G/O_p(F)$  of prime order  $q$ . Denote by  $S$  the preimage of  $\langle s \rangle$  in  $G$ . We need to show that  $S \leq \langle \text{tor}(G) \rangle$ . If not then  $S$  is torsion free and therefore is projective by Serre's result ([S-1965]). Then  $q \neq p$  because otherwise  $O_p(F) = O_p(S)$  and so the maximal pro- $p$  quotient  $S/O_p(S) = \langle s \rangle$  has to be projective (cf. Proposition 7.6.7 in [RZ-2000]). So  $q \neq p$  and therefore the maximal pro- $p$  quotient of  $S$  is trivial. Hence  $S \leq \langle \text{tor}(G) \rangle$ .

Thus by factoring out the kernel of the epimorphism of  $F$  to its maximal pro- $p$  quotient we may assume that  $F$  is free pro- $p$ . Let  $M$  be the normal subgroup of  $G$  generated by all elements of order coprime to  $p$ . For a subset  $B$  of  $G$  write  $\bar{B} := BM/M$  and put  $L := FM$ .

**Claim.**  $L/\langle \text{tor}(L) \rangle$  is free pro- $p$ .

Consider the 5-term Hochschild-Serre sequence

$$\begin{aligned} 0 \longrightarrow H^1(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p) &\xrightarrow{\text{Inf}} H^1(\bar{L}, \mathbf{F}_p) \longrightarrow \\ H^1(\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p)^{\bar{L}/\langle \text{tor}(\bar{L}) \rangle} &\xrightarrow{\text{tr}} H^2(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p) \xrightarrow{\text{Inf}} H^2(\bar{L}, \mathbf{F}_p) \end{aligned}$$

By Corollary 2.3  $\text{tor}(\bar{L}) = \overline{\text{tor}(L)}$ . So it suffices to show that  $H^2(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p) = 0$ .

We first show that

$$H^2(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p) \xrightarrow{\text{Inf}} H^2(\bar{L}, \mathbf{F}_p)$$

is 0-map. Denote by  $S_p$  a Sylow subgroup of  $L$  and consider the following commutative diagram:

$$H^2(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p)[r] \xrightarrow{(0.6)} \text{Inf}[d] H^2(\bar{L}, \mathbf{F}_p)[d] H^2(S_p/\langle \text{tor}(S_p) \rangle, \mathbf{F}_p)[r] \xrightarrow{(0.6)} \text{Inf} H^2(S_p, \mathbf{F}_p)$$

where the vertical maps are induced by the natural homomorphisms  $S_p \longrightarrow \bar{L}$  and

$$S_p/\langle \text{tor}(S_p) \rangle \longrightarrow L/\langle \text{tor}(L) \rangle = \bar{L}/\langle \text{tor}(\bar{L}) \rangle.$$

By Theorem 1.7 and Lemma 2.1 the lower horizontal map is 0-map. Since  $M$  is generated by elements of order coprime to  $p$  one has

$$H^1(M, \mathbf{F}_p)^{L/M} = \text{Hom}(M, \mathbf{F}_p)^{(L/M)} = 0.$$

So the 5-term Hochschild-Serre sequence relating cohomology of  $L$  and  $\bar{L}$  implies that

$$\text{Inf} : H^2(\bar{L}, \mathbf{F}_p) \longrightarrow H^2(L, \mathbf{F}_p)$$

is injective. On the other hand

$$\text{Res} : H^2(L, \mathbf{F}_p) \longrightarrow H^2(S_p, \mathbf{F}_p)$$

is injective as well (cf. Lemma 10.2.1 in [W-1998]). Hence the right vertical map is injective. So the commutativity of the diagram implies that

$$H^2(\bar{L}/\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p) \xrightarrow{\text{Inf}} H^2(\bar{L}, \mathbf{F}_p)$$

is 0-map.

Now it suffices to show that

$$H^1(\bar{L}, \mathbf{F}_p) \longrightarrow H^1(\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p)_{\bar{L}/\langle \text{tor}(\bar{L}) \rangle}$$

is surjective. We show that the dual map

$$H_1(\langle \text{tor}(\bar{L}) \rangle, \mathbf{F}_p)_{\bar{L}/\langle \text{tor}(\bar{L}) \rangle} \longrightarrow H_1(\bar{L}, \mathbf{F}_p)$$

is injective. Note that the map in question coincides with the natural homomorphism  $\langle \text{tor}(\bar{L}) \rangle / (\Phi(\langle \text{tor}(\bar{L}) \rangle) [\langle \text{tor}(\bar{L}) \rangle, \bar{L}]) \longrightarrow \bar{L} / \Phi(\bar{L})$ . Pick  $\bar{g} \in \langle \text{tor}(\bar{L}) \rangle \cap \Phi(\bar{L})$ . We need to show that  $\bar{g} \in \Phi(\langle \text{tor}(\bar{L}) \rangle) [\langle \text{tor}(\bar{L}) \rangle, \bar{L}]$ .

As was observed above every torsion element of  $\bar{L}$  lifts to some torsion element of  $L$  and since  $M$  is generated by all elements of  $p'$ -order, it also lifts to an element of  $p$ -power order. Since all the Sylow subgroups of  $L$  are conjugate this implies that  $\text{tor}(\bar{L}) = \text{tor}(S_p)$ ; indeed if  $s \in L$  is element of  $p$ -power order, then  $s^k \in S_p$  for some  $k \in L$  and so  $\bar{s}^k = \bar{s}^l$  for some  $l \in L$  and one can choose an element  $l \in S_p$  such that  $\bar{s}^{kl^{-1}} = \bar{s}$ . It follows that  $\bar{g}$  has a preimage in  $\langle \text{tor}(S_p) \rangle$  (i.e. lifts to an element  $g \in \langle \text{tor}(S_p) \rangle$ ). Since  $\bar{g} \in \Phi(\bar{L})$ ,  $g \in L^p[L, L]M$ . As  $\bar{L}$  is pro- $p$ , one has  $\Phi(S_p)M = L^p[L, L]M$  and so  $g \in \Phi(S_p)M$ . Write  $g = sm$  for some  $s \in \Phi(S_p)$ ,  $m \in M \cap S_p$ .

Put  $\tilde{L} := L / (\Phi(F) \cap M)$  and for  $A \subseteq L$  write  $\tilde{A}$  for the image of  $A$  in  $\tilde{L}$ . Observe that  $\Phi(\tilde{F}) = \Phi(\tilde{F})$ . Then  $\tilde{F} \cap \tilde{M}$  is an elementary abelian pro- $p$  normal subgroup of  $\tilde{F}$ . Moreover, it is central in  $\tilde{F}$  since  $\tilde{M} \cap \Phi(\tilde{F}) = 1$ . Then  $\tilde{F} = B \times (\tilde{F} \cap \tilde{M})$ , where  $B$  is the preimage in  $\tilde{F}$  of the direct complement of  $(\tilde{F} \cap \tilde{M})\Phi(\tilde{F})/\Phi(\tilde{F})$  in  $\tilde{F}/\Phi(\tilde{F})$ . It follows that  $\tilde{L} = \tilde{F}\tilde{M} = B \times \tilde{M}$ , so  $\tilde{S}_p = B \times (\tilde{M} \cap \tilde{S}_p)$ . Then  $\Phi(\tilde{S}_p) = \Phi(B) \times \Phi(\tilde{M} \cap \tilde{S}_p)$  and so we can write  $\tilde{s} = \tilde{s}_0\tilde{m}_0$ , where  $\tilde{s}_0 \in \Phi(B)$ ,  $\tilde{m}_0 \in \Phi(\tilde{M} \cap \tilde{S}_p)$ . Then  $\tilde{g} = \tilde{s}_0\tilde{m}_0\tilde{m}$ . Note that  $\langle \text{tor}(\tilde{S}_p) \rangle = (\tilde{M} \cap \tilde{S}_p) \times \langle \text{tor}(B) \rangle$ , so  $\tilde{s}_0 \in \langle \text{tor}(B) \rangle$ . Since  $\Phi(F) \cap M \leq \Phi(S_p)$ , using again the equality  $\text{tor}(\bar{L}) = \text{tor}(S_p)$  one infers that there exist  $s_0 \in \Phi(S_p) \cap \langle \text{tor}(S_p) \rangle$ ,  $m' \in M$  such that  $g = s_0m'$ . But  $S_p$  is virtually free pro- $p$  by Theorem 1.7, so  $H_2(S_p/\langle \text{tor}(S_p) \rangle, \mathbf{F}_p) = 0$ . Hence from 5-term Hochschild-Serre exact sequence relating  $S_p$  and  $S_p/\langle \text{tor}(S_p) \rangle$  one concludes that

$$H_1(\langle \text{tor}(S_p) \rangle, \mathbf{F}_p)_{S_p/\langle \text{tor}(S_p) \rangle} \longrightarrow H_1(S_p, \mathbf{F}_p)$$

is injective or equivalently

$$\langle \text{tor}(S_p) \rangle / \Phi(\langle \text{tor}(S_p) \rangle)[\langle \text{tor}(S_p) \rangle, S_p] \longrightarrow S_p / \Phi(S_p)$$

is injective. This means that

$$\Phi(S_p) \cap \langle \text{tor}(S_p) \rangle = \Phi(\langle \text{tor}(S_p) \rangle)[\langle \text{tor}(S_p) \rangle, S_p]$$

and so

$$s_0 \in \Phi(\langle \text{tor}(S_p) \rangle)[\langle \text{tor}(S_p) \rangle, S_p].$$

Therefore  $\bar{g} \in \Phi(\langle \text{tor}(\bar{L}) \rangle)[\langle \text{tor}(\bar{F}) \rangle, \bar{L}]$  and the claim is proved.

Now, the preceding claim shows that  $\bar{G} = G/M$  is virtually free pro- $p$ , and since by Corollary 2.3  $G/\langle \text{tor}(G) \rangle = \bar{G}/\langle \text{tor}(\bar{G}) \rangle$ , the result follows from Theorem 1.7.  $\square$

**Theorem 2.5.** *Let  $G$  be a virtually projective group. Then  $G/\langle \text{tor}(G) \rangle$  is projective.*

*Proof:* It suffices to show that for a Sylow subgroup  $S_p$  of  $G$  one has  $S_p \langle \text{tor}(G) \rangle / \langle \text{tor}(G) \rangle$  is free pro- $p$  for every  $p$ . However, this follows from Lemma 2.4 since  $S_p \langle \langle \text{tor}(G) \rangle / \langle \text{tor}(G) \rangle \rangle$  is a pro- $p$  group.  $\square$

### 3. Examples

We conclude this section with two examples: an example of a pro-2 group  $G$  with  $\text{vcd}(G) = 2$  such that  $G/\langle \text{tor}(G) \rangle$  contains torsion and an example of a pro-2 group  $H$  containing a free pro-2 subgroup  $F$  of index 2 that does not satisfy the conclusion of Theorem 1.2.

The first example shows that groups of virtual cohomological dimension 1 are exceptional with respect to the property studied in the paper.

The second example shows that the existence of a section  $\mathcal{T}/G \longrightarrow G$  is essential for Theorem 1.2.

**Example.** Let  $G = \mathbf{Z}_2 \amalg_H D_\infty$ , where  $D_\infty$  is the infinite dihedral pro-2 group and  $H$  is the subgroup of order 2 in both factors (note that  $D_\infty$  contains a unique subgroup of index 2 isomorphic to  $\mathbf{Z}_2$ ). Then the normal closure  $N$  of the first factor is of index 2 in  $G$  and isomorphic to the generalized dihedral group  $\mathbf{Z}_2 \rtimes \mathbf{Z}_2$  where the action is by inversion. Hence  $\text{cd}(N) = 2$ . However, the group  $\langle \text{tor}(G) \rangle$  is the normal closure of the second factor and  $G/\langle \text{tor}(G) \rangle$  has order 2 and so is not torsion free.

Before constructing the second example we need the following

**Lemma 3.1.** *Let  $Z \cong \mathbf{Z}_2$ . There exists a profinite  $Z$ -space  $X$  with one point fixed, all other points having trivial stabilizers and the natural surjection  $\pi_Z : X \rightarrow X/Z$  does not admit a continuous section.*

*Proof:* Let  $X$  be a direct product of uncountably many copies of  $\mathbf{Z}_2$ . Define an action of  $Z$  on  $X$  by coordinatewise multiplication. Then the trivial element of  $X$  is the unique fixed point and the stabilizers of all other points of  $X$  are trivial. By Lemma in [CP-1992]  $\pi$  does not admit a continuous section.  $\square$

Now we start to construct the group  $H$ . Let  $(X, Z)$  be as in Lemma 3.1 and  $\mathcal{H}_0 = X \times C$ , where  $C \cong C_2$ . We are going to use the definition of a free pro-2 product in sense of [M-1989] (see also [NSW-2000, Chapter IV, § 3]. Denote by  $(\mathcal{H}_0, pr_X, X)$  the associated constant sheaf. Put  $H_0 = \coprod_X \mathcal{H}_0$ . Define an action of  $Z$  on  $\mathcal{H}_0$  by setting  $(x, c)z = (xz, c)$ ,  $x \in X$ ,  $c \in C$ . Then the universal property of the free pro-2 product allows one to extend this action canonically to a continuous action of  $Z$  on  $H_0$ . Put  $H = H_0 \rtimes Z$ . Using the canonical morphism  $\omega : \mathcal{H}_0 \rightarrow H$ , define  $C_x = \omega(\mathcal{H}_0(x)) \cong C_2$ , and regard  $H_0$  as the internal free pro-2 product  $H_0 = \coprod_{x \in X} C_x$  (see (1.16) and (1.17) in [M-1989] or Chapter IV, § 3 in [NSW-2000]). Denote by  $x_0$  a point which is fixed by  $Z$ . Let  $F_0$  be the free subgroup of index 2 in  $H_0$ . The subset  $W = (\{c_{x_0}^{-1}c_x \mid x \in X\})$  is clearly a closed subset of  $F_0$  and, in fact, is a pointed basis of  $F_0$ . Indeed,  $H_0/\Phi(F_0) = F_0/\Phi(F_0) \times C_{x_0}\Phi(F_0)/\Phi(F_0)$  from where it follows that  $W$  generates  $F_0$ . On the other hand, if  $Y$  is a proper closed subset of  $(\{c_{x_0}^{-1}c_x \mid x \in X\})$ , then  $\overline{\langle Y, C_{x_0} \rangle} = \coprod_{y \in Y} \langle c_{x_0}y \rangle \neq H_0$ , so  $Y$  can not generate  $F_0$ . Observe that  $Z$  acts on  $W$  as follows:  $(c_{x_0}^{-1}c_x)^z = c_{x_0}^{-1}(c_x)^z = c_{x_0}^{-1}c_{xz}$ .

**Lemma 3.2.**

- (a) For  $x_1, x_2 \in X$  one has that  $C_{x_1}$  is conjugate in  $H$  to  $C_{x_2}$  if and only if  $x_1z = x_2$  for some  $z \in Z$ ;
- (b)  $H$  contains a free pro-2 subgroup  $F$  of index 2;

*Proof:* (a) Suppose  $C_{x_1}^h = C_{x_2}$  for some  $h \in H$ . Let  $h_0 \in H_0$ ,  $z \in Z$  be such that  $h = zh_0$ . Then  $C_{x_1}^{zh_0} = (C_{x_1z})^{h_0} = C_{x_2}$ , whence we have  $x_1z = x_2$ , as required. Conversely, if  $x_1z = x_2$  then there exists  $h_0 \in H_0$  such that  $c_{x_1} = c_{x_1z}^{h_0} = c_{x_1}^{zh_0}$  as needed.

(b) Put  $F = F_0 \rtimes Z$ . Let  $f : F_0 \rightarrow F(W/Z)$  be the natural epimorphism induced by the natural surjection  $W \rightarrow W/Z$ . Since  $F(W/Z)$  is free,  $f$  splits, i.e., there exists a monomorphism  $\varphi : F(W/Z) \rightarrow F_0$  with  $f\varphi = \text{id}$ . It suffices to show that the natural homomorphism  $F(W/Z) \amalg Z \rightarrow F$  induced by  $\varphi$  and by the monomorphism sending  $Z$  to its copy in  $F$  is an isomorphism. But this is clear since this homomorphism induces an isomorphism on the Frattini quotients (cf. Theorem 7.2.7 in [RZ-2000]).  $\square$

**Proposition 3.3.**  *$H$  cannot be isomorphic to a free product of centralizers of finite groups and a free pro-2 group.*

*Proof:* Suppose there exists a boolean space  $T$ , and a continuous family  $\Sigma_H := \{C_t \mid t \in T\}$  of groups of order 2 such that  $H = \coprod_{t \in T} C_H(C_t) \amalg L$  for some free pro-2 group  $L$ . Note that  $\coprod_{t \in T} C_t$  is a subgroup of  $H_0$ . For  $t \in T$  denote by  $c_t$  the generator of  $C_t$  and put  $S_T = \{c_t \mid t \in T\}$ . Similarly for  $x \in X$  denote by  $c_x$  the generator of  $C_x$  and put  $S_X = \{c_x \mid x \in X\}$ . Let  $\bar{S}_T$  and  $\bar{S}_X$  be the homeomorphic images of  $S_T$  and  $S_X$  in  $H/\Phi(H_0)$  respectively. By Proposition 4.9 in [M-1989] every finite subgroup of  $H$  is

conjugate to one of its factor  $C_x$ . Therefore,  $\bar{S}_T \subseteq \bar{S}_X$ . Let  $f : H_0/\Phi(H_0) \longrightarrow H/\Phi(H)$  be the natural homomorphism. Since all  $C_t$  are subgroups of free factors  $C_H(C_t)$  of  $H$ , the restriction  $f|_{S_T}$  is an injection. It is also a surjection because any finite subgroup is conjugate to a subgroup of a free factor (see Proposition 4.9 in [M-1989]). Now observe that the  $Z$ -set  $S_X$  is isomorphic to the  $Z$ -set  $\bar{S}_X$ , where abusing notation we use the same letter for the image of  $Z$  in  $H/\Phi(H_0)$  and, by Lemma 3.2 (a), it is isomorphic to the  $Z$ -set  $X$  as well. Also note that the restriction  $f|_{\bar{S}_X}$  coincides with the natural quotient map  $\bar{S}_X \longrightarrow \bar{S}_X/Z$ . On the other hand  $f|_{S_T} : \bar{S}_T \longrightarrow \bar{S}_X/Z$  is also a surjection by Lemma 3.2 (a) taking into account that every  $C_x$  is conjugate to some  $C_t$  in  $H$  (see Proposition 4.9 in [M-1989]). Thus  $f|_{S_T} : \bar{S}_T \longrightarrow \bar{S}_X/Z$  is a homeomorphism. Since, as was mentioned above,  $Z$ -set  $\bar{S}_X$  is isomorphic to the  $Z$ -set  $X$ , we obtain a contradiction with the fact established in Lemma 3.1 that a continuous section  $X/Z \longrightarrow X$  does not exist.  $\square$

## REFERENCES

- [CP-1992] Z. A. Chatzidakis and P. Pappas [1992]. Von Neumann regular rings not representable as rings of continuous functions, *Algebra Universalis*, **29**, 332–337.
- [DS-1975] J.L. Dyer and G.P. Scott [1975] Periodic automorphisms of free groups. *Comm. in Algebra* **3** (1975) 195-201.
- [HRZ-1999] W.N Herfort, L. Ribes and P.A. Zalesskii.  $p$  - Extensions of Free Pro- $p$  Groups. *Forum Math.* **11** (1998), 49-61.
- [HZ-2002] W.N Herfort and P.A. Zalesskii. Virtually Free pro- $p$  Groups. Preprint.
- [M-1989] O.V. Melnikov. Subgroups and Homology of Free Products of Profinite Groups. *Math. USSR Izvestiya*, **34**, 1, (1990), 97-119.
- [NSW-2000] J. Neukirch, A. Schmidt and K. Wingberg, Cohomology of Number Fields, Springer 2000.
- [RZ-2000] L. Ribes and P.A. Zalesskii, Profinite Groups. Springer 2000.
- [RZ1-2000] L. Ribes and P.A. Zalesskii, Pro- $p$  Trees. (2000), Chapter 3, Ser. Progress in Mathematics, Birkhäuser Boston. Ed. M. du Sautoy, A. Shalev, D. Segal.
- [Sch-1994] C. Scheiderer [1994]. *Real and Étale Cohomology*, Lect. Notes Math., **1588**, Springer, Berlin.
- [Sch-1994] C. Scheiderer [1994]. The structure of some virtually free pro- $p$  groups. *Proc. Amer. Math. Soc.* **127** (1999) 695-700.
- [S-1965] J-P. Serre, Sur la dimension cohomologique des groupes profinis, *Topology* **3** (1965) 413-420.
- [W-1998] J.S. Wilson. Profinite Groups. London Math.Soc. Monographs (Clarendon Press, Oxford, 1998)
- [ZM-1990] P.A. Zalesskii and O.V. Melnikov. Fundamental Groups of Graphs of Profinite Groups. *Algebra i Analiz*, **1** (1989); translated in: *Leningrad Math. J.* **1** (1990), 921-940.

Departamento de Matemática  
Universidade de Brasília  
70910-900 Brasília-DF  
Brasil